

In the Claims:

Please cancel claims 31, 32, 33, and 37.

Please amend the claims as follows:

14. (Amended) A compound according to claim 6 wherein X is O.

20. (Amended) A compound according to claim 4 wherein

R¹ is phenyl which is substituted in the *meta* position with one or more substituents selected from the group consisting of halogen, -CF₃, C₁₋₈alkyl, C₁₋₈alkylamino, alkoxy, C₃₋₆cycloalkylC₂₋₆alkenyl, C₆₋₁₄arylC₂₋₆alkenyl, -CN, -NO₂, -NH₂, -SR⁶, -S(O)₂R⁶, -S(O)R⁷, -S(O)₂R⁷, -C(O)R⁷, C₂₋₆alkenyl which may be optionally substituted with a substituent selected from the group consisting of hydroxy, halogen, aryl, and heterocycle, and C₂₋₆alkynyl which may be optionally substituted with a substituent selected from the group consisting of hydroxy, halogen, aryl, C₃₋₆cycloalkyl, and heterocycle;

R² is hydrogen;

R³ is hydrogen;

R⁴ is phenyl substituted in the *ortho* position with a substituent selected from the group consisting of hydroxy, halogen, -CF₃, or C₁₋₈alkyl and substituted at the *para* position with a substituent selected from the group consisting of hydroxy, halogen, -CF₃, C₁₋₈alkyl, hydroxyC₁₋₈alkyl, -CN, -NO₂, C₁₋₈alkylamino, heterocycleC₁₋₈alkyl, -C(O)NH₂, -S(O)R⁷, -S(O)₂R⁷, -C(O)R⁷, -NS(O)₂R⁷, -S(O)₂NR⁸R⁹, -S(O)₂NHR¹¹, -SO₂R¹¹, -OR¹¹, -C(O)R¹¹, -C(O)NR¹¹, -C(O)OR¹¹, -NR¹¹, -NC(O)R¹¹, heterocycleC₂₋₆alkenyl, heterocycle which may be optionally substituted with one or more substituents selected from the group consisting of oxo, C₁₋₈alkyl, and C(O)OR¹¹, and C₁₋₈alkyl which may be optionally substituted with one or more substituents selected from the group consisting of -CN and heterocycle, optionally substituted with -C(O)R¹¹;

R⁵ is a substituent in the *para* position relative to X and is selected from the group consisting of halogen, C₁₋₈alkyl, -NO₂, -NH₂, C₁₋₈alkylamino, CF₃, or alkoxy;

a3 R¹¹ is C₁₋₈alkyl, optionally substituted with one or more substituents selected from the group consisting of hydrogen, C₁₋₈alkyl, -S(O)₂NR⁸R⁹, -NR⁸R⁹, and heterocycle, optionally substituted with one or more substituents selected from the group consisting of oxo and C₁₋₈alkyl; or a pharmaceutically acceptable derivative thereof.

a4 26. (Amended) A compound according to claim 4 wherein R¹ is C₆₋₁₄ aryl substituted in the meta position, particularly with halogen and wherein R³ is hydrogen and R⁴ is C₆₋₁₄aryl substituted with C₁₋₈alkyl, in particular methyl.

27. (Amended) A method of treatment of a viral infection in a mammal comprising administering to said mammal an antivirally effective amount of a compound according to claim 2.

a5 29. (Amended) A method of inhibiting HIV reverse transcriptase comprising administering to a mammal an effective amount of a compound according to claim 2.

30. (Amended) A method of preventing HIV infection, or of treating HIV infection, comprising administering to a mammal an effective amount of a compound according to claim 2.

a6 34. (Amended) A pharmaceutical composition comprising an effective amount of a compound according to claim 2 together with a pharmaceutically acceptable carrier.

Please add the following new claims:

38. A compound according to claim 6 wherein

a7 R¹ is phenyl which is substituted in the *meta* position with one or more substituents selected from the group consisting of halogen, -CF₃, C₁₋₈alkyl, C₁₋₈alkylamino, alkoxy, C₃₋₆cycloalkylC₂₋₆alkenyl, C₆₋₁₄arylC₂₋₆alkenyl, -CN, -NO₂, -NH₂, -SR⁶, -S(O)₂R⁶, -S(O)R⁷, -S(O)₂R⁷, -C(O)R⁷, C₂₋₆alkenyl which may be optionally substituted with a substituent selected from the group consisting of hydroxy, halogen, aryl, and heterocycle, and C₂₋₆alkynyl which may be optionally

substituted with a substituent selected from the group consisting of hydroxy, halogen, aryl, C₃₋₆cycloalkyl, and heterocycle;

R² is hydrogen;

R³ is hydrogen;

67
R⁴ is phenyl substituted in the *ortho* position with a substituent selected from the group consisting of hydroxy, halogen, -CF₃, or C₁₋₈alkyl and substituted at the *para* position with a substituent selected from the group consisting of hydroxy, halogen, -CF₃, C₁₋₈alkyl, hydroxyC₁₋₈alkyl, -CN, -NO₂, C₁₋₈alkylamino, heterocycleC₁₋₈alkyl, -C(O)NH₂, -S(O)R⁷, -S(O)₂R⁷, -C(O)R⁷, -NS(O)₂R⁷, -S(O)₂NR⁸R⁹, -S(O)₂NHR¹¹, -SO₂R¹¹, -OR¹¹, -C(O)R¹¹, -C(O)NR¹¹, -C(O)OR¹¹, -NR¹¹, -NC(O)R¹¹, heterocycleC₂₋₆alkenyl, heterocycle which may be optionally substituted with one or more substituents selected from the group consisting of oxo, C₁₋₈alkyl, and C(O)OR¹¹, and C₁₋₈alkyl which may be optionally substituted with one or more substituents selected from the group consisting of -CN and heterocycle, optionally substituted with -C(O)R¹¹;

R⁵ is a substituent in the *para* position relative to X and is selected from the group consisting of halogen, C₁₋₈alkyl, -NO₂, -NH₂, C₁₋₈alkylamino, CF₃, or alkoxy;

R¹¹ is C₁₋₈alkyl, optionally substituted with one or more substituents selected from the group consisting of hydrogen, C₁₋₈alkyl, -S(O)₂NR⁸R⁹, -NR⁸R⁹, and heterocycle, optionally substituted with one or more substituents selected from the group consisting of oxo and C₁₋₈alkyl; or a pharmaceutically acceptable derivative thereof.

39. A compound according to claim 7 wherein

R¹ is phenyl which is substituted in the *meta* position with one or more substituents selected from the group consisting of halogen, -CF₃, C₁₋₈alkyl, C₁₋₈alkylamino, alkoxy, C₃₋₆cycloalkylC₂₋₆alkenyl, C₆₋₁₄arylC₂₋₆alkenyl, -CN, -NO₂, -NH₂, -SR⁶, -S(O)₂R⁶, -S(O)R⁷, -S(O)₂R⁷, -C(O)R⁷, C₂₋₆alkenyl which may be optionally substituted with a substituent selected from the group consisting of hydroxy, halogen, aryl, and heterocycle, and C₂₋₆alkynyl which may be optionally substituted with a substituent selected from the group consisting of hydroxy, halogen, aryl, C₃₋₆cycloalkyl, and heterocycle;

R² is hydrogen;

R³ is hydrogen;

a7
R⁴ is phenyl substituted in the *ortho* position with a substituent selected from the group consisting of hydroxy, halogen, -CF₃, or C₁₋₈alkyl and substituted at the *para* position with a substituent selected from the group consisting of hydroxy, halogen, -CF₃, C₁₋₈alkyl, hydroxyC₁₋₈alkyl, -CN, -NO₂, C₁₋₈alkylamino, heterocycleC₁₋₈alkyl, -C(O)NH₂, -S(O)R⁷, -S(O)₂R⁷, -C(O)R⁷, -NS(O)₂R⁷, -S(O)₂NR⁸R⁹, -S(O)₂NHR¹¹, -SO₂R¹¹, -OR¹¹, -C(O)R¹¹, -C(O)NR¹¹, -C(O)OR¹¹, -NR¹¹, -NC(O)R¹¹, heterocycleC₂₋₆alkenyl, heterocycle which may be optionally substituted with one or more substituents selected from the group consisting of oxo, C₁₋₈alkyl, and C(O)OR¹¹, and C₁₋₈alkyl which may be optionally substituted with one or more substituents selected from the group consisting of -CN and heterocycle, optionally substituted with -C(O)R¹¹;

R⁵ is a substituent in the *para* position relative to X and is selected from the group consisting of halogen, C₁₋₈alkyl, -NO₂, -NH₂, C₁₋₈alkylamino, CF₃, or alkoxy;

R¹¹ is C₁₋₈alkyl, optionally substituted with one or more substituents selected from the group consisting of hydrogen, C₁₋₈alkyl, -S(O)₂NR⁸R⁹, -NR⁸R⁹, and heterocycle, optionally substituted with one or more substituents selected from the group consisting of oxo and C₁₋₈alkyl; or a pharmaceutically acceptable derivative thereof.

40. A compound according to claim 17 wherein

R¹ is phenyl which is substituted in the *meta* position with one or more substituents selected from the group consisting of halogen, -CF₃, C₁₋₈alkyl, C₁₋₈alkylamino, alkoxy, C₃₋₆cycloalkylC₂₋₆alkenyl, C₆₋₁₄arylC₂₋₆alkenyl, -CN, -NO₂, -NH₂, -SR⁶, -S(O)₂R⁶, -S(O)R⁷, -S(O)₂R⁷, -C(O)R⁷, C₂₋₆alkenyl which may be optionally substituted with a substituent selected from the group consisting of hydroxy, halogen, aryl, and heterocycle, and C₂₋₆alkynyl which may be optionally substituted with a substituent selected from the group consisting of hydroxy, halogen, aryl, C₃₋₆cycloalkyl, and heterocycle;

R² is hydrogen;

R³ is hydrogen;

an R^4 is phenyl substituted in the *ortho* position with a substituent selected from the group consisting of hydroxy, halogen, $-CF_3$, or C_{1-8} alkyl and substituted at the *para* position with a substituent selected from the group consisting of hydroxy, halogen, $-CF_3$, C_{1-8} alkyl, hydroxy C_{1-8} alkyl, $-CN$, $-NO_2$, C_{1-8} alkylamino, heterocycle C_{1-8} alkyl, $-C(O)NH_2$, $-S(O)R^7$, $-S(O)_2R^7$, $-C(O)R^7$, $-NS(O)_2R^7$, $-S(O)_2NR^8R^9$, $-S(O)_2NHR^{11}$, $-SO_2R^{11}$, $-OR^{11}$, $-C(O)R^{11}$, $-C(O)NR^{11}$, $-C(O)OR^{11}$, $-NR^{11}$, $-NC(O)R^{11}$, heterocycle C_{2-6} alkenyl, heterocycle which may be optionally substituted with one or more substituents selected from the group consisting of oxo, C_{1-8} alkyl, and $C(O)OR^{11}$, and C_{1-8} alkyl which may be optionally substituted with one or more substituents selected from the group consisting of $-CN$ and heterocycle, optionally substituted with $-C(O)R^{11}$;

R^5 is a substituent in the *para* position relative to X and is selected from the group consisting of halogen, C_{1-8} alkyl, $-NO_2$, $-NH_2$, C_{1-8} alkylamino, CF_3 , or alkoxy;

R^{11} is C_{1-8} alkyl, optionally substituted with one or more substituents selected from the group consisting of hydrogen, C_{1-8} alkyl, $-S(O)_2NR^8R^9$, $-NR^8R^9$, and heterocycle, optionally substituted with one or more substituents selected from the group consisting of oxo and C_{1-8} alkyl; or a pharmaceutically acceptable derivative thereof.

41. A compound according to claim 18 wherein

R^1 is phenyl which is substituted in the *meta* position with one or more substituents selected from the group consisting of halogen, $-CF_3$, C_{1-8} alkyl, C_{1-8} alkylamino, alkoxy, C_{3-6} cycloalkyl C_{2-6} alkenyl, C_{6-14} aryl C_{2-6} alkenyl, $-CN$, $-NO_2$, $-NH_2$, $-SR^6$, $-S(O)_2R^6$, $-S(O)R^7$, $-S(O)_2R^7$, $-C(O)R^7$, C_{2-6} alkenyl which may be optionally substituted with a substituent selected from the group consisting of hydroxy, halogen, aryl, and heterocycle, and C_{2-6} alkynyl which may be optionally substituted with a substituent selected from the group consisting of hydroxy, halogen, aryl, C_{3-6} cycloalkyl, and heterocycle;

R^2 is hydrogen;

R^3 is hydrogen;

R^4 is phenyl substituted in the *ortho* position with a substituent selected from the group consisting of hydroxy, halogen, $-CF_3$, or C_{1-8} alkyl and substituted at the

para position with a substituent selected from the group consisting of hydroxy, halogen, $-\text{CF}_3$, C_{1-8} alkyl, hydroxy C_{1-8} alkyl, $-\text{CN}$, $-\text{NO}_2$, C_{1-8} alkylamino, heterocycle C_{1-8} alkyl, $-\text{C}(\text{O})\text{NH}_2$, $-\text{S}(\text{O})\text{R}^7$, $-\text{S}(\text{O})_2\text{R}^7$, $-\text{C}(\text{O})\text{R}^7$, $-\text{NS}(\text{O})_2\text{R}^7$, $-\text{S}(\text{O})_2\text{NR}^8\text{R}^9$, $-\text{S}(\text{O})_2\text{NHR}^{11}$, $-\text{SO}_2\text{R}^{11}$, $-\text{OR}^{11}$, $-\text{C}(\text{O})\text{R}^{11}$, $-\text{C}(\text{O})\text{NR}^{11}$, $-\text{C}(\text{O})\text{OR}^{11}$, $-\text{NR}^{11}$, $-\text{NC}(\text{O})\text{R}^{11}$, heterocycle C_{2-6} alkenyl, heterocycle which may be optionally substituted with one or more substituents selected from the group consisting of oxo, C_{1-8} alkyl, and $\text{C}(\text{O})\text{OR}^{11}$, and C_{1-8} alkyl which may be optionally substituted with one or more substituents selected from the group consisting of $-\text{CN}$ and heterocycle, optionally substituted with $-\text{C}(\text{O})\text{R}^{11}$;

R^5 is a substituent in the *para* position relative to X and is selected from the group consisting of halogen, C_{1-8} alkyl, $-\text{NO}_2$, $-\text{NH}_2$, C_{1-8} alkylamino, CF_3 , or alkoxy;

R^{11} is C_{1-8} alkyl, optionally substituted with one or more substituents selected from the group consisting of hydrogen, C_{1-8} alkyl, $-\text{S}(\text{O})_2\text{NR}^8\text{R}^9$, $-\text{NR}^8\text{R}^9$, and heterocycle, optionally substituted with one or more substituents selected from the group consisting of oxo and C_{1-8} alkyl; or a pharmaceutically acceptable derivative thereof.

42. A compound according to claim 19 wherein

R^1 is phenyl which is substituted in the *meta* position with one or more substituents selected from the group consisting of halogen, $-\text{CF}_3$, C_{1-8} alkyl, C_{1-8} alkylamino, alkoxy, C_{3-6} cycloalkyl C_{2-6} alkenyl, C_{6-14} aryl C_{2-6} alkenyl, $-\text{CN}$, $-\text{NO}_2$, $-\text{NH}_2$, $-\text{SR}^6$, $-\text{S}(\text{O})_2\text{R}^6$, $-\text{S}(\text{O})\text{R}^7$, $-\text{S}(\text{O})_2\text{R}^7$, $-\text{C}(\text{O})\text{R}^7$, C_{2-6} alkenyl which may be optionally substituted with a substituent selected from the group consisting of hydroxy, halogen, aryl, and heterocycle, and C_{2-6} alkynyl which may be optionally substituted with a substituent selected from the group consisting of hydroxy, halogen, aryl, C_{3-6} cycloalkyl, and heterocycle;

R^2 is hydrogen;

R^3 is hydrogen;

R^4 is phenyl substituted in the *ortho* position with a substituent selected from the group consisting of hydroxy, halogen, $-\text{CF}_3$, or C_{1-8} alkyl and substituted at the *para* position with a substituent selected from the group consisting of hydroxy, halogen, $-\text{CF}_3$, C_{1-8} alkyl, hydroxy C_{1-8} alkyl, $-\text{CN}$, $-\text{NO}_2$, C_{1-8} alkylamino,

an heterocycleC₁₋₈alkyl, -C(O)NH₂, -S(O)R⁷, -S(O)₂R⁷, -C(O)R⁷, -NS(O)₂R⁷, -S(O)₂NR⁸R⁹, -S(O)₂NHR¹¹, -SO₂R¹¹, -OR¹¹, -C(O)R¹¹, -C(O)NR¹¹, -C(O)OR¹¹, -NR¹¹, -NC(O)R¹¹, heterocycleC₂₋₆alkenyl, heterocycle which may be optionally substituted with one or more substituents selected from the group consisting of oxo, C₁₋₈alkyl, and C(O)OR¹¹, and C₁₋₈alkyl which may be optionally substituted with one or more substituents selected from the group consisting of -CN and heterocycle, optionally substituted with -C(O)R¹¹;

R⁵ is a substituent in the *para* position relative to X and is selected from the group consisting of halogen, C₁₋₈alkyl, -NO₂, -NH₂, C₁₋₈alkylamino, CF₃, or alkoxy;

R¹¹ is C₁₋₈alkyl, optionally substituted with one or more substituents selected from the group consisting of hydrogen, C₁₋₈alkyl, -S(O)₂NR⁸R⁹, -NR⁸R⁹, and heterocycle, optionally substituted with one or more substituents selected from the group consisting of oxo and C₁₋₈alkyl; or a pharmaceutically acceptable derivative thereof.

43. A compound according to claim 6 wherein R¹ is C₆₋₁₄ aryl substituted in the meta position, particularly with halogen and wherein R³ is hydrogen and R⁴ is C₆₋₁₄aryl substituted with C₁₋₈alkyl, in particular methyl.

44. A compound according to claim 7 wherein R¹ is C₆₋₁₄ aryl substituted in the meta position, particularly with halogen and wherein R³ is hydrogen and R⁴ is C₆₋₁₄aryl substituted with C₁₋₈alkyl, in particular methyl.

45. A compound according to claim 17 wherein R¹ is C₆₋₁₄ aryl substituted in the meta position, particularly with halogen and wherein R³ is hydrogen and R⁴ is C₆₋₁₄aryl substituted with C₁₋₈alkyl, in particular methyl.

46. A compound according to claim 18 wherein R¹ is C₆₋₁₄ aryl substituted in the meta position, particularly with halogen and wherein R³ is hydrogen and R⁴ is C₆₋₁₄aryl substituted with C₁₋₈alkyl, in particular methyl.

47. A compound according to claim 19 wherein R¹ is C₆₋₁₄ aryl substituted in the meta position, particularly with halogen and wherein R³ is hydrogen and R⁴ is C₆₋₁₄ aryl substituted with C₁₋₈ alkyl, in particular methyl.

48. A method of treatment of a viral infection in a mammal comprising administering to said mammal an antivirally effective amount of a compound according to claim 4.

an 49. A method of treatment of a viral infection in a mammal comprising administering to said mammal an antivirally effective amount of a compound according to claim 23.

50. A method of inhibiting HIV reverse transcriptase comprising administering to a mammal an effective amount of a compound according to claim 4.

51. A method of inhibiting HIV reverse transcriptase comprising administering to a mammal an effective amount of a compound according to claim 23.

52. A method of preventing HIV infection, or of treating HIV infection, comprising administering to a mammal an effective amount of a compound according to claim 4.

53. A method of preventing HIV infection, or of treating HIV infection, comprising administering to a mammal an effective amount of a compound according to claim 23.

54. A pharmaceutical composition comprising an effective amount of a compound according to claim 4 together with a pharmaceutically acceptable carrier.

55. A pharmaceutical composition comprising an effective amount of a compound according to claim 23 together with a pharmaceutically acceptable carrier.